

TRINITY LABAN CONSERVATOIRE OF MUSIC & DANCE

Research Online at Trinity Laban

Associations between static and dynamic field balance tests in assessing postural stability of female undergraduate dancers

Clarke, F., Koutedakis, Y., Wilson, M., Wyon, M.

Journal of Dance Medicine & Science 25(3)

1 **Associations Between Static and Dynamic Field Balance Tests in Assessing Postural**
2 **Stability of Female Undergraduate Dancers**

3 Frances Clarke, PhD^{1,2}, Yiannis Koutedakis, PhD^{3,4}, Margaret Wilson, PhD⁵, Matthew
4 Wyon, PhD^{2,4}

5
6 ¹Trinity Laban Conservatoire of Music & Dance, London, UK

7 ²National Institute of Dance Medicine and Science, UK

8 ³Department of Sport and Exercise Sciences, University of Thessaly, Trikala, Greece

9 ⁴Sport and Physical Activity Research Centre and School of Performing Arts, University of
10 Wolverhampton, Walsall, UK

11 ⁵Department of Theatre and Dance, University of Wyoming, Laramie, Wyoming, USA

12

13 **Conflict of Disclosure:** None

14

15 **Correspondence address:**

16 Frances Clarke
17 Faculty of Dance,
18 Trinity Laban Conservatoire of Music & Dance,
19 Creekside,
20 London, SE8 3DZ
21 UK

22

23 Phone: +44 020 8305 9422

24 Email: f.clarke@trinitylaban.ac.uk

25

26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51

Abstract

Balance testing on dancers has revealed a wide variety of assessment tools. However, as most field balance tests have been developed for either sport or elderly populations, the evidence of associations between tests and their functional relevance to dance is inconclusive. We assessed possible associations between five such field balance tests. The total of 83 female undergraduate dance students (20 ± 1.5 years; 163 ± 6.6 cm; 61 ± 10.8 kg) volunteered for the tests. They executed the Star Excursion Balance Test (SEBT), the modified Romberg test, the Airplane test, the BioSway Balance System (Biodex, USA) and a dance-specific pirouette test. Spearman's correlation coefficients examined relationships between the measures of the balance tests. Results showed quite strong to strong relationships between some SEBT reach directions ($p < 0.01$), and very weak to moderate relationships between some balance tests including some SEBT directions, Romberg, Airplane, Biosway, and pirouette ($p < 0.01$ and $p < 0.05$). Our findings suggest that current tests used to assess dancers' postural stability need further investigation to ensure functionality and relevance.

Introduction

Field static and dynamic balance tests are useful tools in assessing dancers' postural stability, as they can be set up and utilised in dance studios and laboratories; they are also quick and efficient to use. The importance of testing balance is widely recognised as important for dancers^{1,2} and an integral part of the assessment of dancers in codified theatrical dance techniques^{2,3} and assessing optimal performance^{4,5}. Field balance tests are frequently used to evaluate postural stability in dance screening programmes at the beginning of a performance or study season³, and/or following injuries and subsequent rehabilitation work^{6,7}. Furthermore, field tests have been utilised in balance studies on dancer-specific skills^{8,9}, comparisons between dancers and athletes¹⁰, and dancers and non-dancers¹¹, and investigations on specific sensory organisation of the visual, proprioceptive and vestibular

52 senses^{3,12}. However, to date, a wide range of field assessment tools and test protocols have
53 been employed for assessing dancers' balance but with no evident replication power^{13,14} or
54 analysis of associations between tests.

55 Balance has been defined as an individual's ability to control equilibrium^{15,16} and is a
56 complex phenomenon in the case of dancers^{3,17}. The balance process maintains the position
57 of the body's centre of gravity over the base of support, relying on continuous, rapid
58 feedback from visual, vestibular, and somatosensory structures and followed by coordinated
59 neuromuscular actions^{18,19}. Balance is required during both locomotion and stance, thus, two
60 major types of balance have been defined for measurement purposes. Static balance is the
61 ability to maintain postural stability with the centre of mass over the base of support with
62 minimal movement or at rest²⁰, whereas dynamic balance is the ability to maintain postural
63 stability with the centre of mass over the base of support with the body in motion²⁰.

64 Theatrical dance genres demand expert skill in both static and dynamic balance.

65 Dancers, like gymnasts, use both quick and slow movements in their repertoire, and
66 often use a small base of support^{13,15,21}. Moreover, many balances in dance relate more to
67 dynamic equilibrium in response to sudden movements such as acceleration, deceleration,
68 and rotation^{16,22,23}. Surprisingly, assessments of ballet dancers' balance ability are based
69 predominately on static balance tests¹³, although assessment measures not utilising force
70 plates, such as field tests, do use more dynamic balance tests²⁴. The majority of static balance
71 tests perform one-legged stance positions^{25,26,27} which may not relate to the complex,
72 dynamic dance movements²⁸ in dance repertoire.

73 As aforementioned, a range of field assessment tools have been utilised to assess
74 dancers' postural stability but the majority of these were developed for sports people and the
75 general population. The Star Excursion Balance Test (SEBT) was originally developed as a
76 rehabilitative tool²⁹ but has been adapted with a number of modifications including the Y

77 Balance Test^{30,31}, and a modified SEBT (m/r SEBT)³². One study which utilised a battery of
78 tests including the SEBT, the Balance Error Scoring System (BESS) and the Modified Bass
79 Test of Dynamic Balance (BASS) found mixed results between dancers and non-dancers'
80 balance ability²⁴. Other field tests have used a bespoke one-legged stance^{8,25}, a modified
81 Romberg test^{33,34}, the Biosway Balance test¹², the Airplane test³⁴, or more complex, dance-
82 specific tasks such as a modified ronds de jambe⁶ and pirouettes^{28,35,36}.

83 Despite the range of studies, and to the best of our knowledge, no previous research
84 has investigated the associations between field balance tests. This limited knowledge in the
85 field may impede the choice of appropriate tests to assess balance ability in dance training,
86 screening and research studies. Therefore, the aim of this study was to assess possible
87 relationships between balance tests assessing static and dynamic balance and to ascertain
88 their relevance³ to measuring dancers' balance. To assess the association between recognised
89 field balance tests, the researchers selected five field tests used in assessing postural stability
90 of adult dancers who were either in full time dance training or working as professional
91 dancers in theatrical dance genres¹⁴. Both static and dynamic balance are essential to dance
92 performance, therefore results were compared between static and dynamic balance tests.
93 Three dynamic balance tests were selected: Star Excursion Balance Test, the pirouette test
94 and the Airplane test, and two static balance tests were selected: modified Romberg and the
95 Biosway test. The tests varied in the nature of their test protocols, which may imply
96 assessment of different aspects of postural stability. However, as the tests selected for this
97 study are commonly used in screening, training programmes, and research tests on dancers,
98 the analysis of possible associations between them was deemed to be important in order to
99 examine their potential functional relevance for dancers. It was hypothesised that there would
100 be no significant relationships between the five field balance tests.

101 **Methods**

102 *Participants*

103 Following approval by a University Ethics Committee, and *a priori* power analysis assuming
104 an 80% power with an alpha level of 5%, a total of 83 female dance undergraduates (age:
105 20 ± 1.5 years; height: 163 ± 6.6 cm; mass: 61 ± 10.8 kg; dance experience: 10.18 ± 2.39 yrs)
106 volunteered for testing. All participants were studying on the same undergraduate dance
107 programme and received equal hours of training in contemporary, ballet and jazz. Inclusion
108 criteria specified that they attended dance classes for a minimum of 8 hours per week, were
109 injury free, and that they were 18 years or older. Prior to testing, participants completed a
110 consent form and a pre-activity health questionnaire and those with a known injury or illness
111 were excluded.

112 *Procedures*

113 Prior to balance testing, anthropometric data were obtained from all volunteers, including leg
114 length. The latter was measured with the participant lying supine, from the anterior superior
115 iliac spine to the medial malleolus using an anthropometric tape measure^{29,38}. Following the
116 initial assessments, all participants completed a 15-minute standardised warm up session. The
117 same researcher conducted the tests and ensured accurate positioning, alignment and
118 performance of all participants during testing. Participants took part in tests in a randomised
119 order; the order of supporting leg was also randomised in each test.

120 *Measures*

121 The Star Excursion Balance Test (SEBT) has shown a strong interrater reliability of
122 $ICC=0.35-0.93$ and intrarater reliability of $ICC=0.78-0.96$ ³⁹. The SEBT is marked out on a
123 grid consisting of 8 lines marked on the floor, extending from a common point at 45° angle
124 increments. The reaching directions were referenced according to the supporting leg as
125 anterior (0°), anteromedial (45°), medial (90°), posteromedial (135°), posterior (180°),
126 posterolateral (225°), lateral (270°), and anterolateral (315°). The test was performed on a

127 single leg stance with the middle of the standing foot over the centre of the grid. The non-
128 weight bearing leg extends along each designated line to maximal reach whilst maintaining
129 the support foot on the floor and an upright posture upright facing the front²⁹ (Figure 1). The
130 SEBT procedure was demonstrated by the researcher and participants performed practice
131 trials to ensure accuracy in alignment and foot placement before the reaching distances were
132 measured. The average of three trials was taken for each leg. The participants were instructed
133 to bend their supporting leg as much as possible and reach in the eight directions, touching
134 the furthest point with the most distal part of the foot. At the point of touchdown of the
135 reaching leg, a mark was made by the researcher. Participants were not allowed to slide the
136 foot or to put weight on the reach foot. Termination of tests criteria were displacement of the
137 supporting foot and if weight was put on the reach foot³⁸. Leg reach distances were measured
138 (cm) for each reach direction from the centre of the grid to the touchdown mark. The reach
139 distances in each direction were normalised to % leg length^{24,29}.



140

141 Figure 1. Participant on the SEBT

142 Performance of the Star Excursion
143 Balance Test using the left leg as the limb
144 stance in the medial direction

145

146 Pirouettes are a recognised dance-specific balance test with en dehors turns being most
147 widely used^{28,35,56}. Although, to date, no pirouette tests have been empirically validated¹⁴,
148 pirouettes are recognised as having functional relevance when measuring dancer's postural
149 stability²⁸. Single en dehors pirouettes^{28,35,36} were selected for this study replicating the
150 predominant use of en dehors pirouettes in published studies¹⁴. In the pirouette test,
151 participants were instructed to perform six single en dehors turns consecutively, starting from
152 and returning to, a small open turned out position of the feet with one foot crossed in front of
153 the other (4th position). Tests were conducted on both legs. The pirouettes were conducted on
154 the ball of the foot (*demi pointe*), and during rotation, both legs were rotated outwards, with
155 the non-weight bearing leg bent with a 90° angle at the knee joint, and toes in contact and
156 placed in front of the knee of the supporting leg (*retiré*). The arms were held in front of the
157 body (1st position) during the rotation. The timing of the sequential turns replicated a
158 commonly used tempo (approximately 96BPM) used in Intermediate level ballet classes, and
159 with which the participants were familiar. Participants wore soft, thin-soled ballet shoes for
160 the pirouette tests. Before testing began, a mark was taped to the floor to signal the start
161 position of the supporting foot. At the start of the test, participants placed the ball (head of the
162 metatarsals) of their front foot on the marker on the floor. At the end of the sixth turn, the
163 final position of the ball of the front foot was marked and the displacement distance from the
164 start mark to the finish mark was measured in centimetres (cm). Termination of tests criteria
165 were the inaccurate placement of feet in the turn preparation position and the non-weight
166 bearing foot touching the floor during a turn.

167 The Airplane test has been determined as a reliable indicator of a dancer's functional balance
168 skill level^{34,40}. The single-leg balance task was conducted in bare feet. The tests started with
169 the non-weight bearing leg extended to the posterior direction creating a horizontal line with
170 the torso which is flexed at 90°. The arms were abducted to 90° in the start position³⁴. The test

171 consisted of five bends of the supporting leg with the arms adducted horizontally in order to
172 touch the floor with the fingertips³⁴. As the support leg extended to return to the start
173 position, the arms abducted horizontally again to 90°. The number of times the fingertips
174 touched the floor was recorded up to, and including, five (0-5) instances. The termination test
175 criterion was displacement of the supporting foot, knee valgus, hip internal rotation, or pelvic
176 drop³⁴.

177 The Romberg test is a widely used neurology test³³ with various modifications^{34,41}. The
178 modified Romberg was selected for this study to provide a potentially greater balance
179 challenge for dancers, replicating an earlier study on dancers³⁴. The test comprised a single-
180 leg balance in a parallel bare foot stance. It was conducted with the non-supporting leg
181 slightly bent and not touching the supporting leg. Arms were crossed across the chest and a
182 blindfold was worn^{34,41}. Romberg tests are commonly measured up to 30 seconds' duration³⁴,
183 subsequently this protocol was followed with the additional data recording of sustained
184 balances up to a minute, so 0-60 seconds, allowing for the participants' healthy profile and
185 skill ability. Termination test criterion was the non-weight bearing foot touching the floor and
186 pronation of the supporting foot.

187 The BioSway™ (Biodex Medical Systems Inc, New York, USA) used for the purposes of
188 this study has shown acceptable intratester reliability of ICC= 0.82-0.43 for stability index
189 and ICC= 0.81-0.55 for foot placement, with the overall stability index scores showing the
190 most reliable stability scores (0.82 for intratester and 0.70 for intertester)⁴². The Biosway
191 Postural Sway test used in this study assessed neuromuscular control by measuring a
192 participant's ability to maintain unilateral postural stability on a static surface using the
193 Stability Index to quantify a participant's ability to maintain their centre of balance in
194 unilateral stance, thus measuring postural sway. The BioSway balance tests were conducted
195 with eyes open in single-leg bare foot stance and participants were asked to look ahead

2196 during the tests. Participants were asked to step onto the platform and to place their arms in a
2197 neutral position. Foot position coordinates marked out on the platform were maintained for
2198 the supporting foot throughout all the trials. Participants performed three 20 second trials on
2199 each leg. Data quantified postural stability: overall stability, anterior/posterior and
2200 medial/lateral, and the overall stability data was recorded for further analysis. Data were
2201 excluded if the non-supporting foot was put down, or if the supporting foot moved from the
2202 marked coordinates.

2203 *Data Analyses*

2204 All variables were tested for normality using the Kolmogorov-Smirnov and Shapiro-Wilk
2205 test. Following the results of testing, Spearman's Rank Order Correlation (ρ) was selected
2206 for correlational analysis of the data. The strength of the value of the correlation coefficient
2207 (ρ) was determined by Cohen's⁴³ guidelines and interpreted based on the following scale:
2208 0.10 to 0.29 (small), 0.30-0.49 (medium), 0.50 to 1.0 (large). Statistical significance was set
2209 at $p < 0.05$ using the SPSS 26 (IBM Corporation, Chicago, Ill).

2210 **Results**

2211 Test descriptive measures are presented in Table 1. Spearman's correlations for all test
2212 variables are presented in Table 2. The strongest correlations were shown for the following
2213 SEBT reach directions: SEBT 45° and SEBT 90° ($r = 0.809$, $p < 0.01$), SEBT 135° and SEBT
2214 180° ($r = 0.808$, $p < 0.01$), SEBT 225° and SEBT 270° ($r = 0.787$, $p < 0.01$), SEBT 0° and
2215 SEBT 45° ($r = 0.776$, $p < 0.01$). Some further fairly strong to moderate correlations between
2216 SEBT reach direction variables can also be seen in Table 2. Otherwise, the Romberg showed
2217 a weak correlation with SEBT 0° ($r = 0.240$, $p < 0.01$), the Pirouette test showed weak
2218 correlations with SEBT 0° ($r = 0.193$, $p < 0.05$), SEBT 45° ($r = 0.202$, $p < 0.05$), SEBT 180°
2219 ($r = -0.203$, $p < 0.05$), SEBT 225° ($r = -0.256$, $p < 0.01$) and SEBT 270° ($r = -0.236$, $p <$
2220 0.01). The BioswayTM showed moderate correlations with SEBT 0° ($r = 0.307$, $p < 0.01$) and

221 SEBT 45° ($r = 0.307$, $p < 0.01$) and weak correlations with SEBT 90° ($r = 0.208$, $p < 0.05$),
 222 SEBT 225° ($r = -0.247$, $p < 0.05$) and SEBT 270° ($r = -0.250$, $p < 0.05$). The Airplane test
 223 showed a weak correlation with the Romberg ($r = 0.295$, $p < 0.01$).
 224

225 Table 1. Mean and Standard Deviation of the measures of the field balance tests
 226

227 <u>Variables</u>	<u>Mean ± SD</u>
SEBT 0° (n=158)	65.53 ± 11.02
SEBT 45° (n=158)	69.31 ± 11.32
SEBT 90° (n=158)	77.10 ± 13.20
SEBT 135° (n=158)	84.86 ± 12.68
SEBT 180° (n=158)	88.39 ± 14.93
SEBT 225° (n=158)	84.07 ± 17.41
SEBT 270° (n=158)	73.14 ± 21.04
SEBT 315° (n=158)	69.12 ± 28.12
Romberg (n=158)	34.55 ± 16.90
Pirouette (n=148)	48.50 ± 31.34
Biosway (n=100)	0.78 ± 0.40
Airplane (n=114)	4.61 ± 0.93

228

229 Note: right and left legs tested so n=total number of leg tests. Units of measurement: SEBT reach directions were measured in centimetres
 230 (cm), Romberg in seconds, Pirouettes in cm, Biosway™ in Stability Index (sway) and Airplane in touches to floor (1-5)

231

232

233

234

235 Table 2 about here (attached at end of paper)

236 (Table 2. Spearman's correlations between measures of field balance tests)

237

238

239

240 **Discussion**

241 The purpose of this study was to assess associations between static and dynamic balance tests
242 used to measure postural stability in dancers. Although our results indicated strong
243 relationships between some SEBT reach directions, other relationships between the balance
244 test variables were weak, except for a moderate correlation between the Biosway and SEBT
245 0° and the Biosway and SEBT 45°. The only correlation not including a SEBT reach direction
246 was between the Airplane and Romberg although this was a weak relationship. In this study
247 the eight SEBT reach directions were assessed rather than a composite SEBT score or the Y
248 test to see if any of the eight directions had an association with each other or with the other
249 balance tests. Those SEBT directions demonstrating the strongest relationships with other
250 directions were close in proximity on the SEBT grid although it is not possible to ascertain
251 potential causes of these associations. In reference to dancers' abilities in the SEBT reach
252 directions, the few studies utilising the SEBT in studies on dance populations have reported
253 mixed results. For example, a randomised controlled trial testing eight SEBT directions
254 following a whole body vibration (WBV) intervention, noted an improvement in the anterior,
255 anteriomedial, medial and anterior lateral directions⁴⁴, whilst in another study, dancers
256 achieved higher scores than non-dancers in the medial and posteriomedial planes of
257 movement²⁴. Currently, there is inconclusive evidence in the literature on dancers' balance
258 ability in the SEBT reach directions.

259 Whilst these five tests have been used previously in research studies on dancers’
260 balance, it was acknowledged that each test has different protocols and conditions, resulting
261 in some variations in assessment of postural stability, and this does not necessarily diminish
262 the value of each task. A key example is the Romberg performed with eyes closed. Mixed
263 findings have been reported on dancers’ balance ability in vision conditions¹⁴ and it has been
264 argued that whilst dance training increases the influence of proprioceptive skills over vision
265 information, dancers’ balance strategies rely on different senses in the multimodal processing
266 depending on the specific balance task⁴⁵. Although clinical assessments have identified
267 classifications of balance and postural control strategies for those with balance problems^{46,47},
268 to date, no such balance tool is available for assessing dancers. The five tests in this study
269 demonstrate some resonance with the clinically based Balance Evaluation Systems Test
270 (BESTest)⁴⁶, most notably, the pirouette in their Anticipatory Postural Adjustments category
271 and the Romberg in their Sensory Orientation category but it should be remembered that the
272 BESTest was designed for a very different population.

273 In previous literature, the SEBT, Airplane, Biosway, Romberg, and Pirouette tests
274 have been identified as reliable or accepted balance tasks for the dance population in previous
275 literature^{9,14,34,35,36,40}. It is possible that, in past research, assumptions have been made about
276 the functionality of the tests for dancers even though there have been clear differences in test
277 conditions, and no replication of studies, for example, pirouette studies which have included a
278 range of differing turn tasks^{9,35,36}. Therefore, the predominately weak associations between
279 these field tests revealed in this study may suggest that some balance measures are inadequate
280 for an accurate assessment of dancers’ postural stability, but this may not diminish the
281 validity for some of the tests for different populations. The participants in this study were
282 undergraduate dancers and injury free and it should be noted that there may be differences in
283 what the tests evaluate for postural stability for alternative populations. For example,

284 different results might be elicited in a symptomatic dance population or for professional
285 dancers.

286 When considering the relevance of balance tests employed in research on dancers,
287 several factors need be considered. To date, screening, research studies, and rehabilitation
288 work with dancers have employed a battery of field balance tests¹⁴ but these tests may have
289 little or no predictive power. The lack of replicated studies in balance research on dancers^{13,14}
290 has implications for the conclusions drawn from balance studies. Assumptions on the
291 functionality and relevance of balance tests for dancers are likely to have been made over the
292 years, but reported results may need to be considered within the context of assessed study
293 limitations in the literature^{14,48,49}.

294 Another factor to be considered when assessing balance tests is the task difficulty.
295 Balance tests do not necessarily produce challenging enough demands for dancers^{50,51,52,53}.
296 Dancers' balance has been found to be more automatized than non-dancers⁵⁰ with greater
297 behavioural flexibility² and less cognitive involvement⁵⁰. They use a wide range of balance
298 strategies to maintain, achieve or restore equilibrium and have fast anticipatory reactions. It
299 has been suggested that dancers may reach a ceiling effect in postural automaticity
300 particularly in eyes open tasks⁵⁰. Further balance study limitations can include levels of
301 expertise⁵⁰, for example, if the task is too simple and not challenging enough for the level of
302 expertise of the dancers being assessed⁵⁴, or alternatively, too demanding⁵⁵. Notwithstanding
303 our results indicating weak correlations between specified static and dynamic balance tests,
304 further investigation in this area of research is recommended.

305 **Strengths and limitations**

306 To our knowledge, this is the first study to examine potential associations between specific
307 balance tests employed to measure dancers' postural stability. The relatively large number of
308 volunteers could also be treated as a study strength⁴⁹. However, the present results may have

309 been subject to certain methodological limitations. There is no agreed definition for the wider
310 construct of postural control or stability for dancers⁴⁷. The postural control and movement
311 complexity required for the SEBT and Airplane could be regarded as only moderately
312 challenging for dancers. In addition, reach distances in the SEBT may have been subjected to
313 participants' own exertion and interpretation of the given instructions. The Biosway may not
314 have posed a sufficient challenge for the participants as it was a static position and resembled
315 a basic element of dance technique. A limitation was that the participants were undergraduate
316 dance students and testing on professional dancers might have yielded different results. Also,
317 there were varying levels of expertise demonstrated in the pirouette test and it is possible that
318 some participants were holding the body in a rigid position due to a learned effect or
319 misperception of the required technique⁵⁵.

320 **Conclusion**

321 The purpose of this study was to investigate the potential associations between static and
322 dynamic balance tests already employed in assessing dancers' postural stability, and to
323 ascertain their relevance for assessing dancers' postural stability. Our findings indicated
324 associations between some SEBT reach directions and certain SEBT directions with the
325 Romberg, Pirouette, and Biosway, and the Airplane and Romberg., Except for the
326 associations between some SEBT directions, the strength of the associations between tests
327 was weak. Overall, these weak associations between tests may suggest that some balance
328 measures have some limitations in assessing accurately dancers' postural stability and may
329 not challenge dancers who have demonstrated greater behavioural flexibility in balance tasks.
330 This study has pointed to the need for further investigation of balance assessment tools
331 utilised to assess dancers' postural stability to help reduce study limitations in this area of
332 research. Furthermore, identification of definitions of the wider construct of postural stability

333 (and postural control) for dancers may enhance the choice and application of measurement
334 tools for dancers in the future.

335 **References**

- 336 1. Hugel F, Cadopi M, Kohler F, Perrin P. Postural control of ballet dancers: a specific
337 use of visual input for artistic purposes. *Int J Sports Med.* 1999;20:86-92.
- 338 2. Schmit JM, Regis DI, Riley MA. Dynamic patterns of postural sway in ballet dancers
339 and track athletes. *Expl Brain Res.* 2005;163(3):370-8. doi: 10.1007/s00221-0042185-
340 6
- 341 3. Batson G. Validating a dance-specific screening test for balance: preliminary results
342 from multisite testing. *Med Probl Perform Art.* 2010;25(3):110-5.
- 343 4. Twitchett E, Angioi M, Koutedakis Y, Wyon M. Video Analysis of Classical Ballet
344 Performance. *J Dance Sci Med.* 2009; 13(4):124-8.
- 345 5. Twitchett EA, Angioi M, Koutedakis Y, Wyon M. Do Increases in Selected Fitness
346 Parameters Affect the Aesthetic Aspects of Classical Ballet Performance? *Med Probl*
347 *Perform Art.* 2011; 26(1):35–8.
- 348 6. Clark T, Redding E. The relationship between postural stability and dancer's past and
349 future lower-limb injuries. *Med Probl Perform Art.* 2012;27(4):197-204.
- 350 7. Allen N, Nevill AM, Brooks JHM, Koutedakis Y, Wyon MA. The effect of a
351 comprehensive injury audit program on injury incidence in ballet: a 3-year
352 prospective study. *Clin J Sport Med.* 2013;23(5):373-8. doi:
353 10.1097/JSM.0b013e3182887f32.
- 354 8. Crotts D, Thompson B, Nahom M, Ryan S, Newton RA. Balance abilities of
355 professional dancers on select balance tests. *J Orthop Sports Phys Ther* 1996;3(1):12-
356 17. doi: 10.2519/jospt.1996.23.1.12
- 357 9. Lin C-W, Chen S-J, Su F-C, Wu H-W, Lin C-F. Differences of ballet turns (pirouette)
358 performance between experienced and novice ballet dancers. *Res Q Exerc Sport.*
359 2014;85:330-40. doi: 10.1080/02701367.2014.930088
- 360 10. Gerbino PG, Griffin ED, Zurawski D. Comparison of standing balance between
361 female collegiate dancers and soccer players. *Gait Posture.* 2007;26:501-507. doi:
362 10.1016/j.gaitpost.2006.11.205.
- 363 11. Krityakiarana W, Jongkamonwiwat N. Comparison of balance performance between
364 Thai classical dancers and non-dancers. *J Dance Med Sci.* 2016;20(2):72-8. doi:
365 10.12678/1089-313X.20.2.72
- 366 12. Rein S, Fabian T, Zwipp H, Rammelt S, Weindel S. Postural control and functional
367 ankle stability in professional and amateur dancers. *Clin Neurophysiol.*
368 2011;122(8):1602-10. doi: 10.1016/j.clinph.2011.01.004
- 369 13. Costa MSS, Ferreira AS, Felicio LR. Static and dynamic balance in ballet dancers: a
370 literature review. *Fisioter Pesq.* 2013;20(3):292-298. doi: 10.1590/S1809-
371 29502013000300016
- 372 14. Clarke F, Koutedakis Y, Wilson M, Wyon M. Balance in theatrical dance
373 performance: A systematic review. *Med Probl Perform Art.* 2018;33(4):276-86.
- 374 15. Grimshaw P, Lees A, Fowler N, Burden A. *Sport and Exercise Biomechanics.*
375 Abingdon, Oxon: Taylor & Francis, 2006.
- 376 16. Hall SB. *Basic Biomechanics*, 5th ed., New York: McGraw-Hill. 2007.

- 377 17. Simmons RW. Sensory organization determinants of postural stability in trained ballet
378 dancers. *Int J Neurosci*. 2005;115:87-97. doi: 10.1080/00207450490512678
- 379 18. Nashner LM. Practical biomechanics and physiology of balance. In: Jacobson GP,
380 Newman CW, Kartush JM(eds): *Handbook of balance function testing*. San Diego,
381 CA: Singular Publishing Group, 1997, pp.261-79.
- 382 19. Pollock AS, Durward BR, Rowe PJ, Paul JP. What is balance? *Clin Rehabil*,
383 2000;14(4):402-6. doi: 10.1191/0269215500cr342oa
- 384 20. Hrysomallis C. Balance ability and athletic performance. *Sports*
385 *Medicine*.2011;41(3):221-32.
- 386 21. Bruyneel AV, Mesure S, Paré JC, Bertrand M. Organization of postural equilibrium in
387 several planes in ballet dancers. *Neurosci Lett*. 2010;485:228-232. doi:
388 10.1016/j.neulet.2010.09.017
- 389 22. Golomer E, Dupui P, Séréni P, Monod H. The contribution of vision in dynamic
390 spontaneous sways of male classical dancers according to student or professional
391 level. *J Physiol*. 1999; 93(3):233-7. doi: 10.1016/S0928-4257(99)80156-9
- 392 23. Tortora GJ, Derrickson B. *Principles of Anatomy and Physiology*, 11th ed., USA: John
393 Wiley & Sons, Inc. 2006.
- 394 24. Ambegaonkar JP, Caswell SV, Winchester JB, Shimokochi Y, Cortes N, Caswell
395 AM. Balance comparisons between female dancers and active nondancers. *Res Q*
396 *Exerc Sport*. 2013;84:24-29. doi: 10.1080/02701367.2013.762287.
- 397 25. Schmitt H, Kuni B, Sabo D. Influence of professional dance training on peak torque
398 and proprioception at the ankle. *Clin J Sport Med*. 2005;15(5):331-9. doi:
399 10.1097/01.jsm.0000181437.41268.56
- 400 26. Golomer E, Mbongo F, Toussaint Y, Cadiou M, Israël I. Right hemisphere in visual
401 regulation of complex equilibrium: the female ballet dancers' experience. *Neurol Res*.
402 2010;32(4):409-415. doi: 10.1179/174313209X382476
- 403 27. Bronner S. Differences in segmental coordination and postural control in a multi-joint
404 dance movement développé arabesque. *J Dance Med Sci*. 2012;16(1):26-35.
- 405 28. Lin C-F, Lee I-J, Liao J-H, Wu H-W, Su F-C. Comparison of postural stability
406 between injured and uninjured ballet dancers. *Am J Sports Med*. 2011;39(6):1324-31.
407 doi: 10.1177/0363546510393943
- 408 29. Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to assess
409 dynamic postural-control deficits and outcomes in lower extremity injury: A literature
410 and systematic review. *J Athl Train*. 2012;47(3):339-57.
- 411 30. Plisky P J, Gorman PP, Butler RJ, Kiesel KB, Underwood FB, Elkins B. The
412 reliability of an instrumented device for measuring components of the star excursion
413 balance test. *N Am J Sports Phys Ther*. 2009;4(2):92-9.
- 414 31. Ambegaonkar JP, Cortes N, Caswell SV, Ambegaonkar GP, Wyon M. Lower
415 extremity hypermobility, but not core muscle endurance influences balance in female
416 collegiate dancers. *Int J Sports Phys Ther*, 2016;11(2):220-229.
- 417 32. Wilson M, Batson G. The m/r SEBT: Development of a Functional Screening Tool
418 for Dance Educators. *Med Probl Perform Art*. 2014;29(4):207-15.
- 419 33. Rogers JH. (1980) Romberg and his test. *J Laryngol Otol*. 1980; 94(12):1401-4. doi:
420 10.1017/S002221510009023X
- 421 34. Richardson M, Liederbach M, Sandow E. Functional criteria for assessing pointe-
422 readiness. *J Dance Med Sci*. 2010;14(3):82-88.
- 423 35. Denardi RA, Ferracioli MC, Rodrigues ST. Informação visual e control postural
424 durante a execução da pirouette no ballet. *Rev Port Cien Desp*. 2008;8(2):241-

- 425 250.[Visual information and postural control during pirouette execution in ballet]
426 Portuguese
- 427 36. Golomer EME, Gravenhorst RM, Toussaint Y. Influence of vision and motor imagery
428 styles on equilibrium control during whole-body rotations. *Somatosens Mot Res.*
429 2009;26(4):105-110. doi: 10.3109/08990220903384968
- 430 37. Mertz L, Docherty C. Self-described differences between legs in ballet dancers. Do
431 they relate to postural stability and ground reaction force measures? *J Dance Med Sci.*
432 2012;16(4):154-60.
- 433 38. Kinzey SJ, Armstrong CW. The reliability of the star-excursion test in assessing
434 dynamic balance. *J Orthop Sports Phys Ther.* 1998;27(5):356-60.
- 435 39. Hertel J, Miller SJ, Denegar CR. Intratester and intertester reliability during the Star
436 Excursion Balance Test. *J Sport Rehabil.* 2000;9(2):104-16.
- 437 40. De Wolf A, McPherson A, Besong K, Hiller C, Docherty C. Quantitative measures
438 utilized in determining pointe readiness in young ballet dancers. *J Dance Med Sci.*
439 2018;22(4):209-217. doi.org/10.12678/1089-313X.22.4.209
- 440 41. Khasnis A, Gokula RM. Romberg's test. *J Postgrad Med.*2003; 49(2):169-72.
- 441 42. Schmitz RJ, Arnold BL. Intertester and intratester reliability of a dynamic balance
442 control using the Biodex Stability System. *J Sport Rehabil.* 1998;7(2):95-101.
- 443 43. Cohen JW. *Statistical power analysis for the behavioral sciences.* (2nd ed).,Hillsdale
444 NJ: Lawrence Erlbaum Associates. 1988
- 445 44. Cloak R, Nevill AM, Clarke F, Day S, Wyon MA. Vibration training improves
446 balance in unstable ankles. *Int J Sports Med.* 2010;31:894-900. doi: 10.1055/s-0030-
447 1265151
- 448 45. Bläsing B, Calvo-Merino B, Cross ES, Jola C, Honisch J, Stevens CJ. Neurocognitive
449 control in dance perception and performance. *Acta Psychol.* 2012;139(2):300-308.
450 doi: 10.1016/j.actpsy.2011.12.005
- 451 46. Horak FB, Wrisley DM, Frank J. The Balance Evaluation Systems Test (BESTest) to
452 differentiate balance deficits. *Phys Ther.* 2009;89(5):484-498. doi:
453 10.2522/ptj.20080071
- 454 47. Dewar R, Claus AP, Tucker K, Johnston LM. Perspectives on postural control
455 dysfunction to inform future research: A Delphi study for children with cerebral
456 palsy. *Arch Phys Med Rehab.* 2017;98(3):463-479.
457 doi.org/10.1016/j.apmr.2016.07.021
- 458 48. Guyatt GH, Oxman AD, Vist G, Kunz R, Brozek J, Alonso-Coello P, Montori V, Akl
459 EA, Djulbegovic B, Falck-Ytter Y, Norris SL, Williams Jr JW, Atkins D, Meerpohl J,
460 Schünemann HJ. GRADE guidelines: 4. Rating the quality of evidence—study
461 limitations (risk of bias). *J Clin Epidemiol* 2011b;64:407-15.
- 462 49. Meader N, King K, Llewellyn A, Norman G, Brown J, Rodgers M, Moe-Byrne T,
463 Higgins JPT, Sowden A, Stewart G. A checklist designed to aid consistency and
464 reproducibility of GRADE assessments: development and pilot validation. *Syst Rev.*
465 2014;3:82. doi: 10.1186/2046-4053-3-82
- 466 50. Stins JF, Michielsen ME, Roerdink M, Beek PJ. Sway regularity reflects attentional
467 involvement in postural control: Effects of expertise, vision and cognition. *Gait*
468 *Posture* 2009;30:106-109.
- 469 51. Burzynska AZ, Finc K, Taylor BK, Knecht AM, Kramer AF. The dancing brain:
470 Structural and functional signatures of expert dance training. *Front Hum Neurosci.*
471 2017;11:1-20. doi: 10.3389/fnhum.2017.00566.

- 472 52. Costa de Mello M, Ferreira AS, Felicio LR. Postural control during different unipodal
473 positions in professional ballet dancers. *J Dance Med Sci.* 2017;21(4):151-155. doi:
474 10.12678/1089-313X.21.4.151
- 475 53. Clarke F, Koutedakis Y, Wilson M, Wyon M. Associations between balance ability
476 and dance performance using field balance tests. *Med Probl Perform Art.*
477 2019;34(3):154-160. doi.org/10.21091/mppa.2019.3026
- 478 54. Lobo da Costa PH, Nora FGSA, Vieira MF, Bosch K, Rosenbaum D. Single leg
479 balancing in ballet: effects of shoe conditions and poses. *Gait Posture.*
480 2013;37(3):419-23
- 481 55. Lott BM, Laws KL. The physics of toppling and regaining balance during a pirouette.
482 *J Dance Med Sci.* 2012;16(4):167-74.
483

484

485

486

487

488

ACCEPTED FOR PUBLICATION

489 Table 2. Spearman's correlation analysis between field balance tests

490

	SEBT 0°	SEBT 45°	SEBT 90°	SEBT 135°	SEBT 180°	SEBT 225°	SEBT 270°	SEBT 315°	Romberg	Pirouette	Biosway	Airplane
SEBT 0°	-	.776**	.600**	.447**	.370**	.205**	.080	.500**	.240**	.193*	.307**	.159
SEBT 45°	-	-	.809**	.569**	.408**	.167*	-.008	.318**	.148	.202*	.300**	.145
SEBT 90°	-	-	-	.728**	.509**	.269**	.030	.256**	.084	.065	.208*	.097
SEBT 135°	-	-	-	-	.808**	.591**	.366**	.506**	.050	-.115	.049	.023
SEBT 180°	-	-	-	-	-	.778**	.549**	.682**	.134	-.203*	-.079	.113
SEBT 225°	-	-	-	-	-	-	.787**	.695**	.065	-.256**	-.247*	.019
SEBT 270°	-	-	-	-	-	-	-	.620**	-.032	-.236**	-.250*	.056
SEBT 315°	-	-	-	-	-	-	-	-	.135	-.074	.030	.164
Romberg	-	-	-	-	-	-	-	-	-	.028	-.092	.295**
Pirouette	-	-	-	-	-	-	-	-	-	-	.100	.033
Biosway	-	-	-	-	-	-	-	-	-	-	-	-.047

Airplane

491

492

SEBT = Star Excursion Balance Test

493

** Correlation is significant at the 0.01 level (2-tailed)

494

* Correlation is significant at the 0.05 level (2-tailed)

495

496

ACCEPTED FOR PUBLICATION